

On the method of estimating emission altitude from relativistic phase shift in pulsars

R. T. Gangadhara

Indian Institute of Astrophysics, Bangalore – 560034, India

ganga@iiap.res.in

ABSTRACT

In the preprint astro-ph/0411161 (hereafter G04b), the term $-(\Omega r/c) \sin \theta' \cos \Theta$ is missing in the equation (13) for $\cos \psi$. So, in this supplementary preprint I give the corrected expression for $\cos \psi$ and the revised derivation of aberration phase shift.

1. PHASE SHIFT OF RADIATION EMITTED BY A PARTICLE (BUNCH)

Consider the particle velocity given by

$$\mathbf{v} = \kappa c \hat{b}_{0t} + \mathbf{v}_{\text{rot}}. \quad (1)$$

By substituting for $\mathbf{v}_{\text{rot}}$ (eq. 3 in G04b) into equation (1) we obtain

$$\mathbf{v} = (\kappa c + \Omega r \sin \theta' \cos \Theta) \hat{b}_{0t} + \Omega r \sin \theta' \sin \Theta \hat{e}_{\perp}. \quad (2)$$

By assuming $|\mathbf{v}| \sim c$, from equation (2) we obtain the parameter

$$\kappa = \sqrt{1 - \left(\frac{\Omega r}{c}\right)^2 \sin^2 \theta' \sin^2 \Theta} - \frac{\Omega r}{c} \sin \theta' \cos \Theta. \quad (3)$$

Using the equations (3) and (4) given in G04b, we can solve for $\hat{e}_{\perp}$ and obtain

$$\hat{e}_{\perp} = \frac{\hat{\Omega} \times \hat{r}}{\sin \theta' \sin \Theta} - \cot \Theta \hat{b}_{0t}. \quad (4)$$

Let ψ be the angle between the rotation axis and $\mathbf{v}$, then we have

$$\begin{aligned} \cos \psi = \hat{\Omega} \cdot \hat{v} &= \cos \zeta \left(\sqrt{1 - \left(\frac{\Omega r}{c}\right)^2 \sin^2 \theta' \sin^2 \Theta} - \frac{\Omega r}{c} \sin \theta' \cos \Theta \right) \\ &= \kappa \cos \zeta, \end{aligned} \quad (5)$$

where $\hat{v} = \mathbf{v}/|\mathbf{v}|$. For $r/r_{\text{LC}} \ll 1$, it reduces to $\psi \sim \zeta$.

1.1. Aberration Angle

If η is the aberration angle, then we have

$$\cos \eta = \hat{b}_{0t} \cdot \hat{v} = \kappa + \frac{\Omega r \sin \theta' \cos \Theta}{c}, \quad (6)$$

$$\sin \eta = \hat{\epsilon}_{\perp} \cdot \hat{v} = \frac{\Omega r}{c} \sin \theta' \sin \Theta. \quad (7)$$

Therefore, from equations (6) and (7), we obtain

$$\tan \eta = \frac{\Omega r}{c} \frac{\sin \theta' \sin \Theta}{\sqrt{1 - (\Omega r/c)^2 \sin^2 \theta' \sin^2 \Theta}}. \quad (8)$$

Hence the radiation beam, which is centered on the direction of $\mathbf{v}$, gets tilted (aberrated) with respect to $\hat{b}_{0t}$ due to rotation.

For $\Omega r/c \ll 1$, it can be approximated as

$$\tan \eta \approx \frac{\Omega r}{c} \sin \theta' \sin \Theta. \quad (9)$$

1.2. Aberration Phase Shift

Consider Figure 5 (G04b) in which ZAD, ZBX, ZCY and DXY are the great circles centered on the neutron star. The small circle ABC is parallel to the equatorial great circle DXY. The unit vector $\hat{b}_{0t}$ represents a field line tangent, which makes the angle ζ with respect to the rotation axis ZO. The velocity unit vector $\hat{v}$ is inclined by the angles η and ψ with respect to $\hat{b}_{0t}$ and ZO, respectively. We resolve the vectors $\hat{b}_{0t}$ and $\hat{v}$ into the components parallel and perpendicular to the rotation axis:

$$\hat{b}_{0t} = \sin \zeta \hat{b}_{0t\perp} + \cos \zeta \hat{\Omega}, \quad (10)$$

$$\hat{v} = \sin \psi \hat{v}_{\perp} + \cos \psi \hat{\Omega}, \quad (11)$$

where the unit vectors $\hat{b}_{0t\perp}$ and $\hat{v}_{\perp}$ lie in the plane of small circle ABC. Next, by solving for $\hat{b}_{0t\perp}$ and $\hat{v}_{\perp}$, we obtain

$$\hat{b}_{0t\perp} = \frac{1}{\sin \zeta} (\hat{b}_{0t} - \cos \zeta \hat{\Omega}), \quad (12)$$

$$\hat{v}_{\perp} = \frac{1}{\sin \psi} (\hat{v} - \cos \psi \hat{\Omega}). \quad (13)$$

By taking scalar product with $\hat{b}_{0t\perp}$ on both sides of equation (13), we obtain

$$\cos(\delta\phi'_{abe}) = \hat{v}_\perp \cdot \hat{b}_{0t\perp} = \frac{1}{\sin\psi}(\hat{v} \cdot \hat{b}_{0t\perp} - \cos\psi \hat{\Omega} \cdot \hat{b}_{0t\perp}). \quad (14)$$

Since $\hat{\Omega}$ and $\hat{b}_{0t\perp}$ are orthogonal, we have

$$\cos(\delta\phi'_{abe}) = \frac{1}{\sin\psi}(\hat{v} \cdot \hat{b}_{0t\perp}). \quad (15)$$

Using $\hat{b}_{0t\perp}$ from equation (12) we obtain

$$\cos(\delta\phi'_{abe}) = \frac{1}{\sin\psi} \frac{(\hat{v} \cdot \hat{b}_{0t} - \cos\zeta \hat{v} \cdot \hat{\Omega})}{\sin\zeta}. \quad (16)$$

By substituting for $\hat{v} \cdot \hat{b}_{0t}$ and $\hat{v} \cdot \hat{\Omega}$ from equations (6) and (5), we obtain

$$\cos(\delta\phi'_{abe}) = \frac{1}{\sin\psi} \frac{(\cos\eta - \cos\zeta \cos\psi)}{\sin\zeta}. \quad (17)$$

Substituting for η again from equation (6), we obtain

$$\cos(\delta\phi'_{abe}) = \frac{1}{\sin\zeta \sin\psi} \left(\kappa + \frac{\Omega r}{c} \sin\theta' \cos\Theta - \cos\zeta \cos\psi \right). \quad (18)$$

By substituting for κ from equation (5), we obtain

$$\cos(\delta\phi'_{abe}) = \frac{1}{\sin\zeta \sin\psi} \left(\frac{\cos\psi}{\cos\zeta} + \frac{\Omega r}{c} \sin\theta' \cos\Theta - \cos\zeta \cos\psi \right). \quad (19)$$

It can be further reduced to

$$\cos(\delta\phi'_{abe}) = \tan\zeta \cot\psi + \frac{\Omega r \sin\theta' \cos\Theta}{c \sin\zeta \sin\psi} \quad (20)$$

Next, by taking a scalar product with $\hat{e}_\perp$ on both sides of equation (13), we obtain

$$\sin(\delta\phi'_{abe}) = \hat{v}_\perp \cdot \hat{e}_\perp = \frac{1}{\sin\psi}(\hat{v} \cdot \hat{e}_\perp - \cos\psi \hat{\Omega} \cdot \hat{e}_\perp). \quad (21)$$

Since $\hat{\Omega} \cdot (\hat{\Omega} \times \hat{r}) = 0$, from equation (4) we obtain

$$\hat{\Omega} \cdot \hat{e}_\perp = -\cot\Theta (\hat{\Omega} \cdot \hat{b}_{0t}). \quad (22)$$

We know from equation (10), $(\hat{\Omega} \cdot \hat{b}_{0t}) = \cos\zeta$. Therefore, we have

$$\hat{\Omega} \cdot \hat{e}_\perp = -\cot\Theta \cos\zeta. \quad (23)$$

By substituting for $\hat{v} \cdot \hat{\epsilon}_\perp$ from equation (7), and for $\hat{\Omega} \cdot \hat{\epsilon}_\perp$ from equation (23) into equation (21) we obtain

$$\begin{aligned} \sin(\delta\phi'_{\text{abe}}) &= \frac{1}{\sin \psi} (\sin \eta + \cos \zeta \cos \psi \cot \Theta) \\ &= \frac{\Omega r \sin \theta' \sin \Theta}{c \sin \psi} + \cos \zeta \cot \psi \cot \Theta . \end{aligned} \quad (24)$$

Thus, we obtain from equations (20) and (24):

$$\tan(\delta\phi'_{\text{abe}}) = \frac{(\Omega r/c) \sin \theta' \sin \Theta + \cos \zeta \cos \psi \cot \Theta}{\tan \zeta \cos \psi + (\Omega r/c)(\sin \theta' \cos \Theta / \sin \zeta)} . \quad (25)$$